

Dynamic Ocean Management: Identifying the Critical Ingredients of Dynamic Approaches to Ocean Resource Management

REBECCA LEWISON, ALISTAIR J. HOBDAI, SARA MAXWELL, ELLIOTT HAZEN, JASON R. HARTOG, DANIEL C. DUNN, DANA BRISCOE, SABRINA FOSSETTE, CATHERINE E. O'KEEFE, MICHELE BARNES, MELANIE ABECASSIS, STEVEN BOGRAD, N. DAVID BETHONEY, HELEN BAILEY, DAVID WILEY, SAMANTHA ANDREWS, LUCIE HAZEN, AND LARRY B. CROWDER

Dynamic ocean management, or management that uses near real-time data to guide the spatial distribution of commercial activities, is an emerging approach to balance ocean resource use and conservation. Employing a wide range of data types, dynamic ocean management can be used to meet multiple objectives—for example, managing target quota, bycatch reduction, and reducing interactions with species of conservation concern. Here, we present several prominent examples of dynamic ocean management that highlight the utility, achievements, challenges, and potential of this approach. Regulatory frameworks and incentive structures, stakeholder participation, and technological applications that align with user capabilities are identified as key ingredients to support successful implementation. By addressing the variability inherent in ocean systems, dynamic ocean management represents a new approach to tackle the pressing challenges of managing a fluid and complex environment.

Keywords: bycatch, sustainable fisheries, stakeholder involvement, remotely sensed data, data integration

The world's oceans are under an unprecedented level of pressure from resource use and commercial activities—for example, fisheries, shipping, aquaculture, and mineral, natural gas, and oil extraction. The impacts from these activities are compounded by climate change, pollution, and invasive species (Halpern et al. 2008, Merrie et al. 2014). In the face of these chronic influences from human activities, ecosystem functions and the services they provide are being eroded (MEA 2005). Ocean management approaches that can address these growing pressures have been the focus of considerable research and agency action (Foley et al. 2010, Ban et al. 2014).

Management of ocean seascapes is complicated by their dynamic spatial and temporal nature. Ocean currents create filaments and eddies that persist for weeks to months (Waugh et al. 2006). Mesoscale variability in ocean structure and processes lead to spatial and temporal patchiness in primary productivity (Hazen et al. 2013). In turn, zooplankton, fish, and other pelagic species in the food chain are mobile and patchily distributed as they seek areas of food to support growth and reproduction (Hauray et al. 1978). As a result, the

ocean is considered highly dynamic in space and time, albeit with persistent physiobiological patterns across a range of scales (e.g., Sheldon et al. 1972, Steele 1980). These dynamic ocean systems are changing—and are predicted to continue to change—in the face of climate change (Polovina et al. 2008, Hazen et al. 2012, Pinsky et al. 2013).

Over the ages, people have often exhibited dynamic behavior in response to changing ocean conditions and have modified or altered their use of ocean resources, particularly fish (Berkes et al. 2000). In this context, dynamic ocean management (DOM) has long been an integral part of human cultures. In the 1400s, fishers in the northwest Atlantic (e.g., Norway, Spain) followed fish populations as local climates changed, resulting in the rise and fall of settlements along the coast (Jackson et al. 2011). Whalers in the early 1900s followed the dynamic ice edge to locate desired species throughout the Arctic (de la Mare et al. 1997). Johannes (1998) describes fishers in modern-day Palau who alter fishing effort on grouper in response to spatial and temporal shifts in spawning aggregation sizes. Several studies have explored shifts in fishing effort in response to changing




Figure 1. Examples of some typical static ocean management structures found worldwide, shown here for Tasmania, Australia: exclusive economic zone (outer blue line), marine protected areas (colored by International Union for Conservation of Nature protection categories—red, IA; blue, II; green, IV; lime, VI), time area closure to protect an inshore squid spawning region (light blue), and a fisheries management zone (grey—trawling closure in waters deeper than 600 meters within the Australian exclusive economic zone).

ocean conditions across many countries, including Solomon Islands, Papua New Guinea, Indonesia, and Hawaii (Aswani 1998, Cinner and McClanahan 2006), and have documented the tactical and strategic decisions fishers make to account for seasonality, changing weather conditions, and levels of bycatch (Christensen and Raakjaer 2006). More recently, ocean use patterns have been shown to change in response to shifting economic conditions. In an analysis of fisher behavior in the French Guyana shrimp fishery, Bene (1996) found that changes in the spatial distribution of fishing effort over time

were more closely associated with market constraints and fluctuating economic incentives than with changes in the spatial distribution of the resource. Scheld and Anderson (2014) describe dynamic landing patterns in which fishers alter concurrent landings of other stocks to avoid flooding markets and dropping market value, a pattern observed in other fisheries (Christensen and Raakjar 2006, Guillen and Maymou 2014).

In contrast to the dynamic nature of ocean organisms, conditions, and users, many ocean management approaches are static (figure 1). Nations have declared exclusive economic

zones (EEZs) to delineate sovereign waters. Ocean management agencies have established fixed marine protected areas (MPAs) and employed time area closures to restrict access and use of areas with sensitive resources. Fisheries managers have set harvest allocations, often at yearly intervals based on synoptic surveys and historical catch information. In many regions these static management constructs have supported resource management, reduced pressure on critical habitats or threatened marine species (Gormley et al. 2012; Pichegru et al. 2010), and in some cases, have resolved conflict among ocean activities or users (Agardy 1994, Gaines et al. 2010). Static management approaches can support management of resource use at fixed locations—for example, offshore energy platforms, or to protect fixed ocean features—for example, coral reefs, but static approaches may be less effective in managing highly mobile organisms, which respond to shifting ocean dynamics and intensifying mobile human pressures (Hyrenbach et al. 2000). Static approaches have failed to prevent fishery closures that are triggered when catch of nontarget species, often referred to as choke species, or protected species exceeds established thresholds (O’Keefe and Decelles 2013). Dynamic ocean management (DOM) is an emerging tool that can support or supplement traditional management approaches for marine ecosystems. However, DOM does not replace management frameworks. Rather, like static approaches, DOM can serve as an instrument to allocate ocean uses and resource extraction within designated thresholds and limits. What differentiates DOM from static management approaches is the use of real-time or near real-time data on the shifting physical, biological, socioeconomic, and other characteristics of the ocean and ocean resource users to generate responsive spatial management measures or strategies (Maxwell et al. 2012, Hobday et al. 2014).

Here, we explore several prominent examples of this emerging management approach to better explain its utility and significance and demonstrate its potential to generate feasible and flexible management actions that can support resources and users. Dynamically managing resource use and commercial activity in marine systems is still fairly uncommon, despite the increasing technical capability to do so (Game et al. 2009, Hobday et al. 2014). We introduce emerging examples of DOM, focus on common threads and critical differences, and evaluate the successes achieved and obstacles faced by DOM approaches. In essence, our review identifies lessons learned in the emerging DOM applications that provide a roadmap to effective DOM development and implementation. Our evaluation summarizes the current state of the science of DOM, articulates the future direction for this promising management tool, and identifies the critical ingredients for DOM success.

Defining and contextualizing dynamic ocean management

Maxwell and colleagues (2012) and Hobday and colleagues (2014) described DOM as management that changes in space and time in response to the shifting nature of the ocean

and its users based on the integration of new biological, oceanographic, social, or economic data. Although DOM is a more recent development in marine resource management, the concepts and potential applications are based on established fields of technical and social scientific endeavor. Information on the dynamic movements of a range of species have been collected with a novel array of biologging devices (Bograd et al. 2010, Block et al. 2011, Hazen et al. 2012), and information on ocean conditions is available from in situ measurements, satellites, and ocean models. There is a wide range of literature describing the creation of habitat models from empirical data sources (Zydelis et al. 2011), to inform the design of MPAs (Bailey and Thompson 2009, McGowan et al. 2013). Similarly, information-sharing networks have been documented for commercial shipping and fishing industries and social science has explored incentives for cooperation amongst ocean resource users (Hardin 1982, Ostrom et al. 1990). Although several types of data and models have been explored, the transfer of these information sources to support management decisions has generally focused on static advice rather than dynamic management approaches (but see Hobday et al. 2010).

Recent advances in the availability, compatibility, and dissemination of long-term biological and remotely sensed environmental data now support the development of dynamic approaches. Likewise, other data types, including social and economic, and genetic information are increasingly accessible and available. Importantly, the ability to disseminate data has rapidly expanded in the last decade with advances in Internet, smart devices, and tablet technologies (Teacher et al. 2013). This rapid paired technology and data expansion supports the defining feature of DOM, which is building management strategies on a range of data layers available in near or real-time. With these data, management products that can support multiobjective or multicriteria management decisions can be developed (Hobday et al. 2014).

Current applications of DOM: Addressing ocean management challenges

DOM approaches are emerging in several places globally, replacing static management approaches to support the management of various mobile ocean activities, including fishing and shipping (table 1). These initiatives have been implemented to maintain target catch within quota limits, reduce bycatch of species of conservation concern, or resolve conflicts among ocean users and uses. Here, we review several examples to illustrate the diversity of applications and the benefits that result from a DOM approach. As the examples demonstrate, the success of these examples derives from innovative combinations and syntheses of complementary environmental, biological, technological, and socioeconomic data.

Maintaining target catch within quota limits. Modern fisheries management is moving towards a system of property rights

Table 1. Key qualitative characteristics and features of the emerging dynamic ocean management (DOM) applications based on literature and expert opinion.

		Yellowtail flounder bycatch avoidance ^a	Turtle Watch ^b	Scottish conserva- tion credits ^c	Pacific Ground- fish fishery ^d	British Columbia salmon fisheries ^e	Eastern Australian longline fishery ^f	Bering Sea Pollock trawl fishery ^g	Whale ALERT ^h	River herring bycatch avoidance ⁱ
Program character- istics	Target quota mgmt			X	X	X				
	Bycatch mgmt	X	X				X		X	
	Avoid protected spp.						X	X		
	Development stage	Oper	Exper	Oper	Oper	Oper	Oper	Oper	Oper	
	Compliance	Vol	Vol	Comp	Vol	Comp	Comp	Vol	Vol	
	Data used	Emp	Mod	Emp	Emp	Emp	Mod	Emp	Emp	
	Type I, II, III, IV	I	III	II	I	I	IV	II	I	
	Time from data received to product release	Within 24 hours	Hourly	1–2 days	Hours	Hours or days	Hours	Hours	Hours	Daily or weekly
Frequency of product update	Daily	Daily	Every 20 days	Within hours	2–3 times per week	Every 2 weeks	4 times per day	Within hours	Daily- weekly	
Data used	Environmental variables		X				X		X	
	Fisheries dependent tags, telemetry, acoustics	X	X	X	X	X		X	X	
	Survey	X								
	Genetic					X				
	Socio-cultural Economic									
Stakeholder involvement	Industry participation (L/M/H)	H	M	H	H	L	L	H	M	H
	Incentives for industry (W/M/S)	S	M	S	M	S	S	S	W	S
	Incentive description	Reduce bycatch, avoid closure	Reduce bycatch, avoid closure	Increased effort allocation	Manage target catch	Manage target catch	Reduce bycatch, avoid closure	Partici- pation linked to higher overall bycatch cap	Avoid collisions	Reduce bycatch, avoid closure
	Partners	Industry University	Govern- ment	Industry Government NGO	Industry NGO	Govern- ment ^s University	Govern- ment ^s	Industry	Govern- ment	Industry Government University
Costs	Cost of program initiation (L/M/H)	L	M	L	M	M	H	M	H	M
	Cost of program maintenance (L/M/H)	L	L	L	M	M	L	M	H	M

Note: The type numbers (Type I-IV) correspond to those in figure 2. The stage of program development was described as operational (Oper) or experimental (Exper). Compliance was described as compulsory (Comp) or voluntary (Vol). Data for programs is either empirical (Emp) or modeled (Mod). Industry participation was described as low (L), moderate (M), or high (H). Incentives for industry were described as weak (W), moderate (M), or strong (S). Relative costs of starting and maintaining DOM programs were categorized as low (L), less than \$100,000; moderate (M), \$100,000 to \$1 million; and high (H), more than \$1 million. Government^s denotes a governmental scientific agency. The references for the programs are the following: ^aO'Keefe and DeCelles 2013. ^bHowell et al. 2008. ^cHolmes et al. 2011, Needle and Catarino 2011. ^dMolteni et al. 2013, www.ecatch.org. ^eBeacham et al. 2004, 2008. ^fHobday et al. 2011. ^gHaflinger and Gruver 2009. ^hWiley et al. 2013, Sibler et al. 2012, <http://stellwagen.noaa.gov/protect/whalealert.html>. ⁱBethoney et al. 2013.

and output controls, often vested in annual quotas that seek to ensure sustainable harvest or promote stock recovery (Hilborn et al. 2003, Gutiérrez et al. 2011). In fisheries operating under quotas, DOM can be used to maintain target catch within allocation limits. Near real-time data can be used to dynamically inform fisheries managers throughout the fishing season allowing them to monitor and manage quota allocations of the target species. For example, British Columbia salmon fisheries are being dynamically managed in order to limit overexploitation of populations of conservation concern, while at the same time supporting the harvest of abundant populations (Beacham et al. 2004, Beacham et al. 2008). The management strategy is based on rapid and highly accurate estimates of stock composition and identification via in-season genetic analyses to allow population-specific exploitation targets to be reached and, generally, quota use to be increased. Similarly, in the New England multispecies groundfish fishery, catch of juveniles from low-quota and unmarketable species (i.e., discards) are counted against the overall fishery quota because juvenile catches degrade stock size and catch of unmarketable species (discards) leads to negative population effects. Dunn and colleagues (2013) provide a theoretical example of DOM that generates rules to guide when a fishing vessel should leave an area, termed a move-on rule. The move-on rules are derived from spatiotemporal autocorrelation analyses that use near real-time data on spatially explicit catch information to reduce unwanted catch of quota-limited and discard species while supporting the harvest of high-quota target species.

Reducing bycatch. Fisheries typically aim to catch one or more target species, but at the same time, can also catch unwanted species, which is termed *bycatch*. Bycatch is a serious issue in many regions, with seabirds, sea turtles, marine mammals, vulnerable sharks, and other fish species impacted (Cullis-Suzuki and Pauly 2010, Croxall et al. 2012, Worm et al. 2013, Lewison et al. 2014). Traditional static management approaches for reducing bycatch include large time-area closures, which can result in serious economic impacts on ocean resource users (Sibert et al. 2012, O’Keefe and Decelles 2013). There are a number of current applications of DOM to reduce bycatch, while still maintaining—and even enhancing—economic efficiency.

One of the best-known DOM examples occurs in the Eastern Australian longline fishery (Hobday and Hartmann 2006; Hobday et al. 2011). Since 2003, a dynamic spatial management approach has been used to limit unwanted capture of southern bluefin tuna (SBT). The approach combines a habitat model conditioned with temperature preference data from pop-up satellite archival tags deployed on SBT, and an ocean model to produce near real-time habitat predictions used by fishery managers during the fishing season. The estimated tuna distribution is used to assist managers in locating management zones that restrict entry by fishers without quota. Three management zones are delineated based on expected SBT distribution and then regulated

through observer coverage and vessel monitoring. The three zones combine the likelihood of tuna presence and the quota status of a fisher. In Zone 1 (OK zone), SBT are not expected to occur and fishers without SBT quota can fish. In Zone 2 (the buffer zone), where there is a limited distribution of SBT predicted, fishers with a limited SBT quota are permitted. In Zone 3 (the core zone), SBT abundance is likely to be highest and only fishers holding sufficient SBT quota may fish and are required to have 100% observer coverage (Hobday et al. 2011).

Another example of the use of DOM to reduce bycatch is TurtleWatch, a tool developed to minimize loggerhead sea turtle (*Caretta caretta*) bycatch in the Hawaii longline swordfish fishery (Howell et al. 2008). Loggerheads are US federally protected species that can be accidentally caught by longline gear. To manage bycatch, an annual loggerhead limit was implemented which, if reached, leads to a fishery-wide closure. Using thermal habitat preferences derived from turtle satellite telemetry data and bycatch occurrences, developers of TurtleWatch found that more than 50% of bycatch interactions occurred between the 17.5 and 18.5 degrees Celsius isotherms. Weekly maps showing the location of both isotherms, areas where loggerhead bycatch is more likely, are computed based on the most recent available sea surface temperature data and distributed online to fishers through the TurtleWatch tool, allowing for voluntary fishing responses. This product has been modified to also include recommended exclusion zones for leatherback turtles (Howell et al. 2015).

On the East Coast of the United States, a yellowtail flounder bycatch avoidance program demonstrates how DOM can be used to reduce bycatch in a scallop fishery. Bycatch levels in relatively small spatial grids (approximately 50 km²) are communicated through vessel monitoring system ship to shore email technology daily or weekly by participating fishing vessels. Researchers summarize the information and provide daily (or weekly) grid-cell specific bycatch advisories to the vessels on the fishing grounds. The program has been used successfully in the Atlantic sea scallop fishery in rotational harvest areas, as well as large regions on Georges Bank and in southern New England to limit yellowtail flounder (*Pleuronectes ferruginea*) bycatch (O’Keefe and DeCelles 2013). Since adopting this bycatch avoidance approach, the fishery has remained open for the designated duration because yellowtail bycatch quota has not been exceeded. This general approach was adapted for the River herring bycatch avoidance program (*Alosa pseudoharengus* and *Alosa aestivalis*) in the US northwest Atlantic midwater and bottom trawl fisheries (Bethoney et al. 2013) with similar success.

The Scottish conservation credits scheme incorporates real-time closures to reduce cod discards and mortality in Scottish and foreign multispecies demersal fisheries. These fisheries target a number of demersal species, including cod (*Gadus morhua*), haddock (*Melanogrammus aeglefinus*) and whiting (*Merluccius merlangus*), but the quotas are set for

each species independently, which leads to significant discards of otherwise marketable fish because they cannot be landed once the quota for that species has been reached. In this DOM application, vessel monitoring systems data and landings are analyzed in gridded fishing areas. On the basis of this analysis, grids with the highest landings per effort are closed to fishing for 21 days if the catch rate of cod exceeds a trigger level. Although evidence of compliance and the direct effects of the closure are difficult to measure, initial analyses of catch savings indicate benefits from the dynamic management regime (Holmes et al. 2011). Grid size and the number of grid closures have been increased to support the goal of limiting discards (Needle and Catarino 2011).

To reduce chinook salmon bycatch, fishers in the Bering Sea pollock fishery are given the opportunity to join bycatch reduction incentive plans which accommodate different processing modes. Participants in the plans are allowed to fish with a higher overall bycatch cap, and all vessels in the fishery have elected to participate. The plans use a rapid response to high bycatch encounters: upon receipt of either an observer report or landing report indicating high salmon bycatch, a short message is sent by email to the entire fleet that contains the coordinates of the encounter and a link to map of the vessel's vessel monitoring system track. The same link allows the user to inspect all high-encounter vessel monitoring system tracks for the week, or for any user-defined period for which data exists on the site. All systems that receive data and generate alerts are fully automated and operate around the clock, so there is almost no delay between receipt of data and the generation of bycatch alerts. On weekly intervals, an analysis of all data received is conducted and fishing areas where bycatch exceeds specified thresholds are closed (Haflinger and Gruver 2009).

Fishers in the Pacific groundfish fishery have also used a DOM approach to reduce bycatch (Molteni 2013). Here, fishers have established risk pools where they share individual transferable fishing quotas for a number of species with low quota, which includes bycatch or traditionally overfished target species. High catch of these species can lead to fishery closures. To avoid capture of overfished target species and the potential of a fishery closure, fishers share data on catch and catch location of these species using a smart-device application called eCatch. Created in partnership with The Nature Conservancy, eCatch collects and shares catch locations of overfished target species in near real-time. Fishers input data on the location and quantity of their overfished target species catch, and a Web-based application serves the data back to fishers in near real-time so they know how much quota of these species have been caught, and what areas to avoid to minimize additional capture of these species.

Resolving direct conflicts. Conflicting management objectives occur in marine sectors other than fisheries, such as shipping, where in some cases species protection goals may be compromised due to deleterious interactions. A smartphone

or tablet application called WhaleALERT (<http://stellwagen.noaa.gov/protect/whalealert.html>) has been developed by resource managers and partners in the Stellwagen Bank National Marine Sanctuary to reduce ship strikes of North Atlantic right whales (*Eubalaena glacialis*), a highly endangered marine mammal with a total population of approximately 522 animals (Pettis and Hamilton 2014). Ships using WhaleALERT receive real-time information on digital nautical charts that includes the location and movement of their ship relative to right whales in the area. The presence of whales is detected by near real-time acoustic buoys that detect right whale calls in the shipping lanes that cross the sanctuary. This information is fed directly into WhaleALERT. Users are alerted when they approach management areas where mandatory or voluntary speed limits are imposed, and compliance is enhanced by pop-up notices that appear on electronic navigation charts when a management boundary is crossed (Wiley et al. 2013). Tools that inform and engage stakeholders may help to improve compliance with mandatory and voluntary measures that are implemented to protect vulnerable species and populations (Silber et al. 2012). Other products developed from whale tracking data, such as seasonal model-based habitat distributions, may also help inform DOM and reduce conflicts between human uses and protected species (Irvine et al. 2014).

Learning from application: Identifying the critical ingredients for DOM success

The emerging applications of DOM provide compelling evidence of the feasibility, efficacy, and utility of the DOM approach. These examples also provide insight into the factors that contribute to DOM success. From a review of the current DOM literature and practice, we identify four key interrelated factors that have influenced the efficacy of DOM approaches: the existing regulatory framework, incentive structure, technological and analytical requirements, and stakeholder participation. More information on the review process can be found in the supplemental material.

Existing regulatory framework and incentive structure. Strong incentives are necessary in order to create successful fisheries management structures (Hilborn et al. 2005, Gutiérrez et al. 2011), a conclusion that also emerges from review of existing DOM approaches (table 1). Existing regulatory frameworks can influence the incentive structure and have been a prominent factor in the establishment of DOM applications. Such regulatory frameworks can include legislation protecting threatened or endangered species (e.g., the US Endangered Species Act or Marine Mammal Protection Act, Australia's Environment Protection and Biodiversity Conservation Act), governing fisheries (e.g., the US Magnuson–Stevens Fishery Conservation and Management Act, the EU Common Fisheries Policy, Australia's Fisheries Management Act), and management directives, such as those through industry management bodies (e.g., US Fishery Management Councils and Commissions). In many of the DOM examples

presented here, the approaches were incentivized or catalyzed directly by the existing regulatory framework, often preceding a large-scale closure or injunction on a particular activity. For example, TurtleWatch was developed to help fishermen avoid reaching a bycatch limit on loggerhead sea turtle bycatch, which if reached can result in a full fishery closure. The yellowtail flounder bycatch avoidance and the Bering Sea pollock trawl fishery programs were developed to help fishers avoid catching nontarget species that had led to complete or partial closures of the fishery with resulting economic loss. Similarly in the Eastern Australian longline fishery, the need to reduce unwanted catch of southern bluefin tuna led to the development of a DOM approach to meet international quota obligations (Hobday et al. 2009). In all three cases, the existing regulatory framework provided the guidelines for the management objectives to be met to maintain fishing activity, and a dynamic approach allowed for an effective means of staying below bycatch quota thresholds, reducing the risk of fishery closures and supporting harvest of target species. Impending regulations can also catalyze the creation of DOM strategies. The development of the River Herring Bycatch Avoidance Program in the US Atlantic herring and mackerel fisheries, was spurred by the proposed creation of river herring protection areas, which would have seasonally closed large portions of the fisheries. Although there may be costs from DOM strategies for ocean resource users, there are also direct and indirect economic and ecological benefits from many of the DOM examples presented here, including increased access to resources, reduced waste and unwanted catch, fewer depleted populations, and more resilient marine assemblages.

Unlike the fisheries examples, the use of dynamic management areas to protect whales from ship collisions with WhaleALERT has a less direct incentive structure for the shipping industry. Vessel collisions are often fatal to the whale and can cause damage to the ship with consequent economic costs. However, these events are rare in comparison to the number of ships at sea and the miles they travel with an average of 11 confirmed or possible ship strikes to large whales per year off the United States based on 292 records between 1975 and 2002 (Jensen and Silber 2004), although this is a minimum as many may go undetected or unreported. To address the threat to whales, vessel speed restrictions have been used as a mitigation measure to reduce the risk of ship strikes for North Atlantic right whales. Compliance with these measures incurs costs by delaying port arrival times, increasing the route length to avoid the management areas or slow down through them. Compliance with the regulations was initially low, but improved with each notification program (Silber et al. 2014). Enforcement activities (citations and fines) appear to be strong motivators and had the greatest influence on improving compliance. Voluntary dynamic measures have been less successful in modifying vessel behavior in this application (Silber et al. 2012). Achieving a balance between economic costs to shipping and benefits to whales, in consultation with

the industry, may allow more effective dynamic management strategies to be developed.

DOM applications have been used primarily in fisheries with a large or established management structure. However, DOM can be relevant for fisheries that do not have robust regulatory structures (i.e., as seen in small-scale fisheries), and industry members have formed partnerships with managers to reduce impacts, even in the absence of regulatory triggers. DOM approaches may help an industry maximize economic gains either by increasing income (e.g., increasing target catch) or minimizing loss (e.g., decreasing bycatch of nuisance species). For example, in small-scale fisheries in Peru, fishers and fisheries observers trade information on unwanted sea turtle bycatch, as well as on ocean and market conditions in real-time using high-frequency two-way radios (Alfaro-Shigueto et al. 2012). The radio program directly benefits the fishers by supplying data on sea turtle bycatch, oceanographic features, and warnings about dangerous conditions. It serves as a vessel safety device, supports economic yield by giving fishers more information on when and where to land their catch, and helps fishers avoid areas of high sea turtle bycatch. Such proactive DOM applications, represent an alternative to reactive approaches to a looming regulatory change, and may support long-term sustainability of a fishery before a crisis develops.

Technological and analytical requirements. Although all DOM approaches require some level of technological and analytical capacity, DOM approaches range from relatively simple approaches that require basic arithmetic to analytical approaches using multiple data streams or complex modeling techniques. Using these examples, it is possible to categorize DOM approaches into four types based on the level of analytical complexity (figure 2). A type I approach involves a single data type and requires minimal data processing. For example, the Yellowtail flounder bycatch avoidance program relies on emailed catch data and creates an easy to follow map as output (type 1). Fishermen send emails to scientists at S Mast with amounts of yellowtail flounder bycatch and target scallop catch by location within grid cell reporting maps that are overlaid on scallop fishing grounds. Scientists then compile data by location and return summarized advisories the next day to highlight areas to avoid in order to minimize yellowtail flounder bycatch. A type II example requires some statistical analysis to integrate near real-time data into a dynamic model. For example, in the theoretical work on the New England multispecies groundfish fishery, the distance a fisher should move from an area to stay below quota limits is determined based on patterns of spatial clustering. A type III approach uses multiple data types to create a dynamic product based on the relationship among input data (e.g., habitat use and environmental data). TurtleWatch relies on more complex data analyses and processing, using location (satellite-tracking or geolocation from archival tags) data and remotely sensed ocean data to model temperature-dependent habitat preferences for loggerhead sea turtles.




Figure 2. The range of complexity of dynamic ocean management (DOM) approaches. In type I, a data stream is processed into a data product for distribution with minimal data processing. In type II, input data are analyzed statistically to produce a dynamic data product. In type III, multiple data types are used to infer statistical relationships, which are then used to build a predictive dynamic model, merged with real-time environmental input, generating data products. In type IV, the output from the dynamic model can be modified by stakeholder adjustments (e.g., manager preferences, additional information on fleet dynamics, compliance considerations) to produce a final data product. Individual data types may be static or can be updated for each run of the model. To date, trade-off analysis has occurred on an informal basis by the stakeholders but could be formally included in future DOM analyses, probably at the type IV end of the spectrum. The dashed lines indicate possible multiple sources of data.

Finally, type IV uses multiple data types and accounts for ocean user behavior. For example, real-time and forecasting habitat models are used in the Eastern Australian longline fishery to regulate fisher access to ocean regions expected to have high numbers of southern bluefin tuna (Hobday and Hartmann 2006, Hobday et al. 2011). In response to these models, fishery managers designate fishing zones that can be accessed depending on fishermen’s available tuna quota. To date, trade-off analysis has occurred on an informal basis by the stakeholders but could be formally included in future DOM analyses (e.g., Abbott 2010).

Although DOM applications can vary in their level of complexity, the technological and analytical tools should align with both the capacity of users to apply the program on the water and the resources available to develop and—critically—maintain the DOM approach. Applying the least labor intensive and most efficient technological and analytical approach will support long-term sustainability of programs by decreasing costs of continued maintenance. Once developed, some analytically complex approaches may be no

more labor intensive than simpler approaches, as computer programming allows for automation of methods (Hobday et al. 2014). For example, the Eastern Australian longline fishery uses a series of automated Matlab scripts to download and integrate remotely sensed data into predefined habitat modeling algorithms (Hartog et al. 2011).

The temporal lag between data capture and management implementation may constrain the types of DOM approaches available to managers. Although WhaleALERT functions in near real-time using acoustic and other data sources, the yellowtail flounder bycatch avoidance program functions on daily time scales, and TurtleWatch uses approximately weekly data. The application of near real-time or time-lagged datasets will further dictate the need for different analytical and technological capacity, both in collecting either remotely sensed or *in situ* type data, and serving it back to users. Shorter time lags may require Internet access on the part of users in remote situations, such as at sea, whereas longer time lags allow for the download of data before departing areas of cell phone or wireless

Internet access. The rapid expansion of smart devices and device-based applications is expanding the options of data sources—for example, crowdsourcing or citizen science, and data delivery (Newman et al. 2012).

Stakeholder participation. Stakeholder collaboration and cooperation have been identified as key components of management success, both in terms of facilitating initial participation and long-term compliance (Österblom et al. 2011). For many of the examples presented here, the development of a DOM application served as a platform to build collaborative networks between scientists, management bodies, industry, and stakeholders (Costanza et al. 1998, Scarlett 2013). Though the parties involved vary among the examples, sometimes including government and nongovernment agencies, the private sector, and academic institutions, all case studies involve some form of partnership with industry and stakeholder participation has been an essential common thread to successful outcomes (see table 1). In the British Columbia salmon fisheries, where a DOM approach has been successful in increasing target species quota use (Beacham et al. 2008), industry was involved from the onset of the program and fishers crowd-sourced information and collected samples for genetic testing. In the yellowtail flounder bycatch avoidance program, which relies on data provided by fishers to successfully meet its management objectives, industry has been involved since the early stages of program development and helped to define the initial program objectives and methods. Extensive collaboration with industry has been similarly fruitful in the Scottish conservation credits cod fishery, and River herring bycatch avoidance program. WhaleALERT was developed with substantial input from shipping companies whose vessels operated in areas subject to regulations protecting right whales, including the development of a test fleet prior to release of the product. In contrast, for TurtleWatch and the Eastern Australian longline fishery, industry were involved primarily as an end user of the products.

Current applications of DOM suggest that building sustained cooperative partnerships among stakeholders, industry, and end users in general in the development of program goals and design may increase the credibility, relevance, legitimacy, and ultimately, the success of DOM. Compliance and participation—whether the program is compulsory or voluntary—can be supported by the establishment of stakeholder networks that work collaboratively to develop and implement the DOM application (O’Keefe and Decelles 2013, Little et al. 2014). Voluntary programs offer an attractive alternative to compulsory ones in many cases because they can be implemented quickly, do not require costly monitoring and penalty regimes, and can promote industry and management innovation (Anton et al. 2004, Silber et al. 2012). However, end-user participation is considered essential in order for voluntary programs to be successful, particularly when monitoring capacity is weak (Bodin et al. 2006, Segerson 2013). In our review, we similarly found strong evidence to suggest that

participation varies depending on the initial level of collaboration with end users, despite the existence of strong incentives (table 1). Even in cases where enforcement capacity is strong, increasing stakeholder collaboration in DOM would likely be advantageous because it can spur self-regulation and curb potential free-riding, thereby reducing enforcement costs (Ostrom 1990, Raakjaer and Mathiesen 2003).

Next steps in dynamic ocean management

DOM approaches are nascent and, as a result, no formal assessment of DOM relative to static management approaches has been conducted. However, our review of existing applications of DOM demonstrates that a suite of management tools have been adopted in multiple sectors across ocean regions, successfully supporting ocean management where competing priorities of economic and ecological outcomes pose challenges. By allowing for explicit incorporation of real-time or near real-time data, these examples show how DOM approaches have the potential to rapidly respond to potential conflicts around ocean resources and support effective management decisionmaking.

One appealing feature of DOM approaches is the ability to provide flexible and fine-scale approaches that balance competing management objectives (figure 3). Traditional static management measures, or those that are updated on seasonal or yearly time scales, encompass comparatively large spatial extents to ensure resource protection. DOM provides an opportunity to create more focused or tailored management structures that can reflect the underlying spatiotemporal relationships between the ocean and the organisms as well as the human activities and the resources being managed. By using real-time or near real-time data, DOM approaches can help to reduce the spatial extent of management actions, fine tuning the management structures and reducing the need for coarse-scale management tactics, even for highly mobile, vagile species.

The current examples of DOM illustrate the utility and relevance of a dynamic management approach and highlight how DOM applications can be coupled with adaptive management in ocean systems as well as the broad goals of ecosystem-based management and marine spatial planning. By directly linking management actions to current conditions, DOM approaches follow the familiar framework of adaptive management—a management structure that supports an iterative decisionmaking and learning process designed to reduce uncertainty in natural system dynamics while concurrently meeting specified management goals and objectives (Walters and Holling 1990, Armitage et al. 2009). DOM approaches align well with the goals of ecosystem-based management as DOM represents place-based management that integrates and recognizes the interdependence across organisms, the environment, and associated processes (biological, oceanographic, social, or economic, *sensu* McLeod et al. 2005). DOM can also support marine spatial planning and zoning efforts. One tool often implemented within the marine spatial planning context is the designation




Figure 3. The relationship between the temporal scale of input data and the spatial extent of management required across a range of species (with variance represented in grey). The species range from a mobile species with a relatively limited distribution (flounder) to a highly vagile species (tuna). Static management measures that are based on data updated on a seasonal or yearly timeframe, shown in black, require comparatively large management areas to ensure resource protection is captured. By using near real-time data on daily, weekly, or monthly time scales, dynamic management, shown in red, allows for the delineation of comparatively smaller areas even for highly vagile species. Abbreviation: km, kilometers.

of marine protected areas (MPAs). In more recent years, calls for dynamic, rather than static, MPAs have intensified to increase MPA effectiveness for highly mobile marine species with variable distributions and in response to projected climate-change induced range shifts (Hyrenbach et al. 2006, Game et al. 2011). Although dynamic MPAs are one potential DOM application, DOM extends far beyond protected areas designation, as the case studies we discuss demonstrate.

The level of complexity with which DOM approaches can be applied is driven by the technological and analytical capacity of both the developers and end users. However, adopting a DOM approach does not necessitate highly complex data input or output as DOM applications have been shown to be successful across a range of complexities (i.e., type I–type IV), as long as the level of complexity of models and the time lags on which data is served to users reflect the ecological and socioeconomic realities of the system being managed. The flexibility and adaptability of this approach presents an appealing feature of DOM; effective DOM approaches can be tailored to meet the needs of the stakeholders. This suggests that DOM can be relevant in both developed and developing country or sector contexts.

Most of the existing DOM examples presented were developed in response to legislative or management action

that threatened to impede or completely curtail ocean uses. Although DOM provided workable solutions in response to these crises, proactive DOM measures are also feasible in balancing the needs of resource users and resources. DOM approaches have also been effective in settings with both voluntary and compulsory compliance regimes. Successful DOM approaches, particularly voluntary or preemptive initiatives, can be facilitated by maintaining active communication channels between resource users, scientists, and managers. This type of ongoing interaction over time also helps to build trust between these different groups, thereby facilitating greater collaboration and cooperation in developing, implementing, and maintaining successful DOM programs. Successful DOM programs that draw from and build on the extensive literature base on stakeholder involvement will likely be most successful in leading to long-term and sustainable change (Gleason et al. 2010, Gopnik et al. 2012).

DOM is an emerging management approach with clear potential, as well as room for improvement and growth. Many of the current DOM examples were designed with a narrow focus on a single species of management concern. As DOM approaches continue to evolve and develop, future DOM applications will likely need to address more complex, multispecies management and include structured trade-off or economic analyses, structured decision-making, and multicriteria decision analysis (White et al. 2012, Klein et al. 2013). There is also a need for future DOM applications to include more social and economic data. In general, stakeholder community structure and diversity (Barnes-Mauthe et al. 2013), attitudes, beliefs, and motivations concerning ocean uses (e.g., Suman et al. 1999), and economic productivity (e.g., Sharma and Leung 1998), have not been explicitly integrated into working applications of DOM. Because communities are often dependent on marine resources to support their livelihoods and humans represent the primary force driving change in marine and coastal systems, lack of integration of these data can directly limit the success of effective management strategies through need for costly enforcement and lack of compliance (see Levin and Lubchenco 2008).

DOM represents an innovative paradigm to tackle the challenges to ocean management and resolve conflicts among multiple objectives by providing an effective alternative to traditional static management. Despite evidence for success in the examples discussed here, there remains a need for

formal evaluations of DOM programs to demonstrate the utility of DOM to industry, management, and other stakeholders. Although no one management approach will resolve the increasing challenge to balance resource use and resource protection, more widespread adoption and implementation of dynamic approaches can play a role in supporting effective ocean management in the twenty-first century.

Acknowledgments

This article was the product of a Center for Ocean Solutions workshop on Dynamic Ocean Management held in December 2013. Funding for this effort was provided by the Center for Ocean Solutions and by NASA Earth Science Division/Applied Sciences Program's ROSES-2012 A.36 Ecological Forecasting (NNH12ZDA001N-ECOF).

Supplemental material

The supplemental material is available online at <http://bioscience.oxfordjournals.org/lookup/suppl/doi:10.1093/biosci/biv018/-/DC1>.

References cited

Agardy MT. 1994. Advances in marine conservation: The role of marine protected areas. *Trends in Ecology and Evolution* 9: 267–270.

Alfaro-Shigueto J, Mangel JC, Dutton PH, Seminoff JA, Godley BJ. 2012. Trading information for conservation: A novel use of radio broadcasting to reduce sea turtle bycatch. *Oryx* 46: 332–339.

Anton WRQ, Deltas G, Khanna M. 2004. Incentives for environmental self-regulation and implications for environmental performance. *Journal of Environmental Economics and Management* 48: 632–654.

Armitage DR, Plummer R, Berkes F, Arthur RI, Charles AT, Davidson-Hunt IJ, Diduck AP, Doubleday NC, Johnson DS, Marschke M et al. 2009. Adaptive co-management for social-ecological complexity. *Frontiers in Ecology and the Environment* 7: 95–102.

Aswani S. 1998. Patterns of marine harvest efforts in southwestern New Georgia, Solomon Islands: Resource management or optimal foraging. *Ocean and Coastal Management* 40: 207–235.

Bailey H, Thompson PM. 2009. Using marine mammal habitat modelling to identify priority conservation zones within a marine protected area. *Marine Ecology Progress Series* 378: 279–287.

Ban NC, Bax NJ, Gjerde KM, Devillers R, Dunn DC, Dunstan PK, Hobday AJ, Maxwell SM, Kaplan DM, Pressey RL 2014b. Systematic conservation planning: A better recipe for managing the high seas for biodiversity conservation and sustainable use. *Conservation Letters* 7: 41–54. doi:10.1111/conl.12010

Barnes-Mauthe M, Arita S, Allen SD, Gray SA Leung PS. 2013. The influence of ethnic diversity on social network structure in a common-pool resource system: Implications for collaborative management. *Ecology and Society*, 18.

Beacham TD, Lapointe M, Candy JR, Miller KM, Withler RE. 2004. DNA in action: Rapid application of DNA variation to sockeye salmon fisheries management. *Conservation Genetics* 5: 411–416.

Beacham TD, Spilsted B, Le KD, Wetklo, M. 2008. Population structure and stock identification of chum salmon (*Oncorhynchus keta*) from British Columbia determined with microsatellite DNA variation. *Canadian Journal of Zoology* 86: 1002–1014.

Béné C. 1996. Effects of market constraints, the remuneration system, and resource dynamics on the spatial distribution of fishing effort. *Canadian Journal of Fisheries and Aquatic Sciences* 53: 563–571.

Bethoney ND, Schondelmeier BP, Stokesbury KDE, Hoffman WS. 2013. Developing a fine scale system to address river herring (*Alosa pseudoharengus*, *A. aestivalis*) and American shad (*A. sapidissima*) bycatch in

the US Northwest Atlantic mid-water trawl fishery. *Fisheries Research* 141: 79–87.

Berkes F, Colding J Folke C. 2000. Rediscovery of traditional ecological knowledge as adaptive management. *Ecological Applications* 10: 1251–1262.

Block BA, et al. 2011. Tracking apex marine predator movements in a dynamic ocean. *Nature* 475: 86–90.

Bodin Ö, Crona B, Ernstson H. 2006. Social networks in natural resource management: What is there to learn from a structural perspective? *Ecology and Society* 11 (resp. 2).

Bograd SJ, Block BA, Costa DP, Godley BJ. 2010. Biologging technologies: New tools for conservation. Introduction. *Endangered Species Research* 10: 1–7.

Christensen A-S, Raakjær J. 2006. Fishermen's tactical and strategic decisions: A case study of Danish demersal fisheries. *Fisheries Research* 81: 258–267.

Cinner JE, McClanahan TR. 2006. Socioeconomic factors that lead to overfishing in small-scale coral reef fisheries of Papua New Guinea. *Environmental Conservation* 33: 73–80.

Costanza R, et al. 1998. Principles for sustainable governance of the oceans. *Science* 281: 198–199.

Croxall JP, Butchart SHM, Laschelles B, Stattersfield AJ, Sullivan B, Symes A, Taylor P. 2012. Seabird conservation status, threats and priority actions: A global assessment. *Bird Conservation International* 22: 1–34.

Cullis-Suzuki S, Pauly D. 2010. Failing the high seas: A global evaluation of regional fisheries management organizations. *Marine Policy* 34: 1036–1042.

De la Mare WK. 1997. Abrupt mid-twentieth-century decline in Antarctic sea-ice extent from whaling records. *Nature* 389: 57–60.

Dunn DC, Boustany AM, Roberts JJ, Brazer E, Sanderson M, Gardner B, Halpin B. 2013. Empirical move-on rules to inform fishing strategies: A New England case study. *Fish and Fisheries* 15: 359–375.

Foley MM, Halpern BS, Micheli F, Armsby MH, Caldwell RM, Crain CM, Prahler E, Rohr N, Sivas D, Beck MW. 2010. Guiding ecological principles for marine spatial planning. *Marine Policy* 34: 955–966.

Gaines SD, White C, Carr MH, Palumbi SR. 2010. Designing marine reserve networks for both conservation and fisheries management. *Proceedings of the National Academy of Sciences* 107: 18286–18293.

Game ET, Grantham HS, Hobday AJ, Pressey RL, Lombard AT, Beckley LE, Gjerde K, Bustamante R, Possingham HP, Richardson AJ. 2009. Pelagic protected areas: The missing dimension in ocean conservation. *Trends in Ecology and Evolution* 24: 360–369.

Gleason M, McCreary S, Miller-Henson M, Ugoretz J, Fox E, Merrifield M, McClintock, W, Serpa P, Hoffman K. 2010. Science-based and stakeholder-driven marine protected area network planning: A successful case study from north central California. *Oceans and Coastal Management* 53: 52–68.

Gopnik M, Fieseler C, Cantral L, McClellan K, Pendleton L, Crowder L. 2012. Coming to the table: Early stakeholder engagement in marine spatial planning. *Marine Policy* 36: 1139–1149.

Gormley AM, Slooten W, Dawson S, Barker RJ, Rayment W, du Fresne S, Brager S. 2012. First evidence that marine protected areas can work for marine mammals. *Journal of Applied Ecology* 49: 474–480. doi:10.1111/j.1365-2664.2012.02121.x

Guillen J, Maynou F. 2014. Importance of temporal and spatial factors in the ex-vessel price formation for red shrimp and management implications. *Marine Policy* 47: 66–70.

Gutiérrez NL, Hilborn R, Defeo O. 2011. Leadership, social capital and incentives promote successful fisheries. *Nature* 470: 386–389.

Haflinger K, Gruver J. 2009. Rolling hot spot closure areas in the Bering Sea Walleye pollock fishery: Estimated reduction of salmon bycatch during the 2006 season. Pages 817–826 in Krueger CC, Zimmerman CE, eds. *Pacific Salmon: Ecology and Management of Western Alaska's Populations*. American Fisheries Society.

Halpern BS, et al. 2008. A global map of human impact on marine ecosystems. *Science* 319: 948–952.

- Hardin R. 1982. Exchange theory on strategic bases. *Social Science Information* 21: 251–272.
- Hartog JR, Hobday AJ, Mearns R, Feng M. 2011. Habitat overlap between southern bluefin tuna and yellowfin tuna in the east coast longline fishery: Implications for present and future spatial management. *Deep-Sea Research Part II: Topical Studies in Oceanography* 58: 746–752.
- Haury LR, McGowan JA, Wiebe PH. 1978. Patterns and processes in the time-space scales of plankton distributions. Pages 277–327 in Steele JH, ed *Spatial Patterns in Plankton Communities*. Plenum Press.
- Hazen EL, Maxwell SM, Bailey H, Bograd SJ, Hamann M, Gaspar P, Godley BJ, Shillinger GL. 2012. Ontogeny in marine tagging and tracking science: Technologies and data gaps. *Marine Ecology Progress Series* 457: 221–240.
- Hazen EL, et al. 2013. Predicted habitat shifts of Pacific top predators in a changing climate. *Nature Climate Change* 3: 234–238.
- Hilborn R, Branch TA, Ernst B, Magnusson A, Minte-Vera CV, Scheuerell MD, Valero JL. 2003. State of the world's fisheries. *Annual Review of Environment and Resources* 28: 359–399.
- Hilborn R, Orensanz JM, Parma AM. 2005. Institutions, incentives and the future of fisheries. *Philosophical Transactions of the Royal Society B* 360: 47–57.
- Hobday AJ, Hartmann K. 2006. Near real-time spatial management based on habitat predictions for a longline bycatch species. *Fisheries Management and Ecology* 13: 365–380.
- Hobday AJ, Hartog, JR, Timmiss T, Fielding J. 2010. Dynamic spatial zoning to manage southern bluefin tuna (*Thunnus maccoyii*) capture in a multi-species longline fishery. *Fisheries Oceanography* 19: 243–253.
- Hobday AJ, Hartog JR, Spillman CM, Alves O. 2011. Seasonal forecasting of tuna habitat for dynamic spatial management. *Canadian Journal of Fisheries and Aquatic Sciences* 68: 898–911.
- Hobday AJ, Maxwell SM, Forgie J, McDonald J, Darby M, Seto K, Bailey H, Bograd SJ, Briscoe DK, Costa DP. 2014. Dynamic ocean management: integrating scientific and technological capacity with law, policy and management. *Stanford Environmental Law Journal* 33: 125–165.
- Hobday AJ, Spillman CM, Hartog JR, Eveson JP. 2015. Seasonal forecasting for decision support in marine fisheries and aquaculture. *Fisheries Oceanography*. doi:10.1111/fog.12083
- Holmes SJ, Bailey N, Campbell N, Catarino R, Barratt K, Gibb A, Fernandes PG. 2011. Using fishery-dependent data to inform the development and operation of a co-management initiative to reduce cod mortality and cut discards. *ICES Journal of Marine Science* 68: 1679–1688.
- Howell EA, Kobayashi DR, Parker DM, Balazs GH, Polovina JJ. 2008 *TurtleWatch*: A tool to aid in the bycatch reduction of loggerhead turtles *Caretta caretta* in the Hawaii-based pelagic longline fishery. *Endangered Species Research* 5: 267–278.
- Howell, EA, Hoover A, Bensen SR, Bailey H, Polovina JJ, Seminoff JA, Dutton PH. 2015. Enhancing the *TurtleWatch* product for leatherback sea turtles, a dynamic habitat model for ecosystem-based management. *Fisheries Oceanography*. 1–12. doi:10.1111/fog.12092
- Hyrenbach KD, Forney KA, Dayton PK. 2000. Marine protected areas and ocean basin management. *Aquatic Conservation: Marine and Freshwater Ecosystems* 10: 437–458.
- Hyrenbach KD, Keiper C, Allen SG, Ainley DG, Anderson DJ. 2006. Use of marine sanctuaries by far-ranging predators: Commuting flights to the California current system by breeding Hawaiian albatrosses. *Fisheries Oceanography* 15: 95–103.
- Irvine LM, Mate BR, Winsor MH, Palacios DM, Bograd SJ, Costa DP, Bailey H. 2014. Spatial and temporal occurrence of blue whales off the U.S. West Coast, with implications for management. *PLOS ONE* (art. e109485). doi:10.1371/journal.pone.0102959
- Jackson JBC, Alexander K, Sala E. 2011. Shifting baselines: The past and the future of ocean fisheries. *Island Press*.
- Jensen AS, Silber GK. 2004. Large whale ship strike database. *National Oceanic and Atmospheric Administration*. Technical Memorandum no. NMFS-OPR-25.
- Johannes, RE. 1998. The case for data-less marine resource management: Examples from tropical nearshore fin fisheries. *Trends in Ecology and Evolution* 13: 243–246.
- Klein CJ, Tulloch, VJ, Halpern BS, Selkoe KA, Watts ME, Steinback C, Scholz A, Possingham HP. 2013 Tradeoffs in marine reserve design: Habitat condition, representation, and socioeconomic costs. *Conservation Letters* 6: 324–332.
- Levin S, Lubchenco J. 2008. Resilience, robustness, and marine ecosystem-based management. *BioScience* 58: 27–32.
- Lewison RL, et al. 2014. Global patterns of marine mammal, seabird, and sea turtle bycatch reveal taxa-specific and cumulative megafauna hotspots. *Proceedings of the National Academy of Sciences* 111: 5271–5276. doi:10.1073/pnas.131896011
- Little AS, Needle CL, Hilborn R, Holland DS, Marshall CT. 2014. Real-time spatial management approaches to reduce bycatch and discards: experiences from Europe and the United States. *Fish and Fisheries*. doi:10.1111/faf.12080
- Maxwell SM, Hazen EL, Morgan LE, Bailey H, Lewison R. 2012. Finding balance in fisheries management. *Science* 336: 413–413.
- McLeod KL, Lubchenco J, Palumbi SR, Rosenberg AA. 2005. Scientific Consensus Statement on Marine Ecosystem-Based Management. *Communication Partnership for Science and the Sea*.
- McGowan J, Hines E, Elliott M, Howar J, Dransfield A, Nur N, Jahncke J. 2013. Using seabird habitat modeling to inform marine spatial planning in Central California's national marine sanctuaries. *PLOS ONE* 8 (art. e71406).
- Merrie A, Dunn DC, Metian M, Boustany AM, Takei Y, Ota Y, Christensen V, Halpin PN, Österblom H. 2014. Human use trends and potential surprise in the global marine common. *Global Environmental Change*. 27: 19–31.
- Millennium Ecosystem Assessment 2005. *Ecosystems and human well-being: Synthesis*. Island Press.
- Molteni M. 2013. How desperation, iPads, and real-time data revived a fishery. *Conservation* (9 September). (31 January 2015; <http://conservationmagazine.org/2013/09/ecatch>)
- Needle CL, Catarino Rui. 2011. Evaluating the effect of real-time closures on cod targeting. *ICES Journal of Marine Science* 68: 1647–1655.
- Newman G, Wiggins A, Crall A, Graham E, Newman S, Crowston K. 2012. The future of citizen science: Emerging technologies and shifting paradigms. *Frontiers in Ecology and the Environment* 10: 298–304.
- O'Keefe CE, DeCelles GR. 2013. Forming a partnership to avoid bycatch. *Fisheries* 38: 434–444.
- Österblom H, Sissenwine M, Symes D, Kadin M, Daw T, Folke C. 2011. Incentives, social-ecological feedbacks and European fisheries. *Marine Policy* 35: 568–574.
- Ostrom E. 1990. *Governing the Commons: The Evolution of Institutions for Collective Action*. Cambridge University Press.
- Pettis H, Hamilton P. 2014. *North Atlantic Right Whale Consortium 2014 Annual Report Card*. North Atlantic Right Whale Consortium.
- Pichegru L, Gremillet D, Crawford RJM, Ryan PG. 2010. Marine no-take zone rapidly benefits endangered penguin. *Biology Letters* 6: 498–501.
- Pinsky ML, Worm B, Fogarty MJ, Sarmiento JL, Levin SA. 2013. Marine taxa track local climate velocities. *Science* 341: 1239–1242.
- Polovina JJ, Howell EA, Abecassis M. 2008. Ocean's least productive waters are expanding. *Geophysical Research Letters* 35 (art. L03618).
- Raakjær Nielsen J, Mathiesen C. 2003. Important factors influencing rule compliance in fisheries lessons from Denmark. *Marine Policy* 27: 409–416.
- Scarlett L. 2013. Collaborative adaptive management: Challenges and opportunities. *Ecology and Society* 18: 26.
- Scheld AM, Anderson CM. 2014. Market effects of catch share management: The case of New England multispecies groundfish. *ICES Journal of Marine Science*. doi:10.1093/icesjms/fsu001
- Segerson K. 2013. Voluntary approaches to environmental protection and resource management. *Annual Review of Resource Economics* 5: 161–180.

- Sharma KR, Leung P. 1998. Technical efficiency of the longline fishery in Hawaii: An application of a stochastic production frontier. *Marine Resource Economics* 13: 259–274.
- Sheldon RW, Prakash A, Sutcliffe WH Jr. 1972. The size distribution of particles in the ocean. *Limnology and Oceanography* 17: 327–340.
- Silber GK, Adams JD, Bettridge S. 2012. Vessel operator response to a voluntary measure for reducing collisions with whales. *Endangered Species Research* 17: 245–254.
- Silber GK, Adams JD, Fonnbeck CJ. 2014. Compliance with vessel speed restrictions to protect North Atlantic right whales. *PeerJ* 2 (art. e399).
- Sibert J, Senina I, Lehodey P, Hampton J. 2012. Shifting from marine reserves to maritime zoning for conservation of Pacific bigeye tuna. *Proceedings of the National Academy of Science* 109: 18221–18225.
- Steele JH. 1980. Patterns in plankton. *Oceanus* 23: 2–8.
- Suman D, Shivlani M, Walter Milon J. 1999. Perceptions and attitudes regarding marine reserves: A comparison of stakeholder groups in the Florida Keys National Marine Sanctuary. *Ocean and Coastal Management* 42: 1019–1040.
- Teacher AGE, Griffiths DJ, Hodgson DJ, Inger R. 2013. Smartphones in ecology and evolution: A guide for the apprehensive. *Ecology and Evolution* 3: 5268–5278.
- Walters C J, Holling C S. 1990. Large-scale management experiments and learning by doing. *Ecology* 71: 2060–2068.
- Waugh DW, Abraham ER, Bowen MM. 2006. Spatial variations of stirring in the surface ocean: A case study of the Tasman Sea. *Journal of Physical Oceanography* 36: 526–542.
- White C, Halpern BS, Kappel CV. 2012. Ecosystem service tradeoff analysis reveals the value of marine spatial planning for multiple ocean uses. *Proceedings of the National Academy of Sciences* 109: 4696–4701.
- Wiley D, Hatch L, Schwehr K, Thompson M, MacDonald C. 2013. Marine sanctuaries and marine planning: Protecting endangered marine life. *United States Coast Guard Proceedings Fall 2013*: 15–20.
- Worm B, Davis B, Kettner L, Ward-Paige CA, Chapman D, Heithaus MR, Kessel ST, Gruber SH. 2013. Global catches, exploitation rates, and rebuilding options for sharks. *Marine Policy* 40: 194–204.
- Zydelis R, et al. 2011. Dynamic habitat models: Using telemetry data to project fisheries bycatch. *Proceedings of the Royal Society B* 278: 3191–3200.

Rebecca Lewison is affiliated with the Department of Biology at Coastal Marine Laboratories, at San Diego State University, in San Diego, California. Alistair J. Hobday, Jason R. Hartog, and Samantha Andrews are affiliated with the Commonwealth Scientific and Industrial Research Organization's Oceans and Atmosphere, in Hobart, Tasmania, Australia. Sara Maxwell is affiliated with the Department of Biological Sciences at Old Dominion University in Norfolk, Virginia. Dana Briscoe and Larry B. Crowder are affiliated with the Stanford University Hopkins Marine Station, in Pacific Grove, California. Elliott Hazen, Sabrina Fossette, and Steven Bograd are affiliated with the National Oceanic and Atmospheric Administration's (NOAA) Southwest Fisheries Science Center's Environmental Research Division, in Pacific Grove. Daniel C. Dunn is affiliated with the Marine Geospatial Ecology Lab at Duke University Marine Lab, in Beaufort, North Carolina. Catherine E. O'Keefe and N. David Bethoney are affiliated with the University of Massachusetts Dartmouth School for Marine Science and Technology, in Fairhaven, Massachusetts. Michele Barnes is affiliated with the Department of Natural Resources and Environmental Management at the University of Hawaii at Manoa, in Honolulu. Melanie Abecassis is affiliated with the Joint Institute for Marine and Atmospheric Research at the University of Hawaii, in Honolulu. Helen Bailey is affiliated with the University of Maryland's Center for Environmental Science's Chesapeake Biological Laboratory, in Solomons, Maryland. David Wiley is affiliated with the NOAA/NOS/Stellwagen Bank National Marine Sanctuary, in Scituate, Massachusetts. Lucie Hazen and LBC are affiliated with the Center for Ocean Solutions, in Monterey, California.